

## A Novel Technique for The Recovery, Isolation and Preliminary Evaluation of *Rhizoctonia solani* Mycoparasites from Soil

Mehmet Hadi AYDIN<sup>1</sup>      Gülay TURHAN<sup>2</sup>

<sup>1</sup> Department of Plant Protection, Faculty of Agriculture, Siirt University, Siirt, Turkey

<sup>2</sup> Seed Technology Application and Research Center (TOTEM), Ege University, İzmir, Turkey

Corresponding Author e-mail: hadiyadin@siirt.edu.tr

Accepted for publication: 14 September 2017

### ABSTRACT

This paper describes a novel technique that allows detecting and isolating of *R. solani* mycoparasites subsist in soil incorporating the precolonized agar method and a modified nutrient medium supplemented with tolcllophos-methyl. A great number of potential biocontrol agents like *Trichoderma* and *Gliocladium* could be isolated into pure culture. Most of the isolates examined macroscopically and microscopically in dual cultures had been recognized as mycoparasites. By using this novel technique, various mycoparasitic isolates were obtained from soil. The majority of the isolates were in the genera of *Trichoderma* and *Gliocladium*. Fourteen *Trichoderma* species, *Gliocladium roseum* Bainier, *Chaetomium* sp., *Geotrichum* sp., *Paecilomyces* sp., *Papulospora* sp. and a *Spicaria* sp. were determined as having mycoparasitic activity against *R. solani* as well. Some of them possessed both parasitic action and antifungal antibiotic activity. This novel technique proved to be useful and showed promise for studies of the ecology of biocontrol agents added to soil or seed.

**Keywords:** Isolation technique, mycoparasite, *Rhizoctonia solani*, *Trichoderma*, *Gliocladium*, tolcllofos methyl

### Topraktaki *Rhizoctonia solani* Mikoparazitlerinin Saptanması, İzolasyonu ve Ön Değerlendirilmesi İçin Yeni Bir Yöntem

### ÖZET

Bu yayın, toprakta bulunan *R. solani* mikoparazitlerini saptamak ve izole etmek amacıyla “petri kabında agarlı ortamda geliştirilmiş *R. solani* kolonisi üzerine toprak örneği ekleyip bir süre beklemek ve bundan cork-borer ile kesilen parçaları tolcllophos-methyl içeren özel besiyerine aktarmak” şeklinde özetlenebilen yeni bir yöntemi açıklamaktadır. Bu yöntemi kullanarak, *Trichoderma* ve *Gliocladium* gibi potansiyel biyokontrol etmenlerinden birçok saf kültür elde edilebilmiştir. İkili kültürlerde makroskopik ve mikroskopik olarak incelenen izolatların çoğunun mikoparazit olduğu anlaşılmıştır. Bu yeni tekniği kullanarak topraktan farklı mikoparazitik izolatlar elde edilmiştir. İzolatların çoğunluğu *Trichoderma* ve *Gliocladium* cinslerine aittir. Ondört farklı *Trichoderma* türü, *Gliocladium roseum* Bainier, *Chaetomium* sp., *Geotrichum* sp., *Paecilomyces* sp., *Papulospora* sp. ve *Spicaria* sp.’nin de *R. solani*’ye karşı mikoparazitik aktiviteye sahip olduğu saptanmıştır. İzolatlardan bazılarının hem parazitik hem de antifungal antibiyotik etkili olduğu anlaşılmıştır. Bu yeni yöntemin toprağa veya tohuma uygulanan biyokontrol etmenlerinin ekolojisi konusundaki çalışmalarda yararlı olacağı öngörülmüştür.

**Anahtar Kelimeler:** İzolasyon yöntemi, mikoparazit, *Rhizoctonia solani*, *Trichoderma*, *Gliocladium*, tolcllofos methyl

### INTRODUCTION

There is an urgent need for new technologies of crop protection. Traditional methods used to protect crops from diseases have been largely based on the use of chemical pesticides. But today there are strict regulations on

A NOVEL TECHNIQUE FOR THE RECOVERY, ISOLATION AND PRELIMINARY EVALUATION OF  
*RHIZOCTONIA SOLANI* MYCOPARASITES FROM SOIL

chemical fungicide use due to carcinogenic effects, residual toxicity problems, environmental pollution and development of fungicide resistant strains but also strong political and public pressure to remove the most hazardous chemicals from the market. Therefore many researchers have focused their efforts on developing alternative inputs to synthetic chemicals for controlling pests and diseases and different strategies have been hypothesized (Pal and Gardener, 2006; Vinale et al., 2008; Abo-Elyousr et al., 2014).

One of the most promising means is the use of new tools based on biocontrol agents. The use of antagonist microorganisms against fungal plant pathogens is an attractive and environmentally-friendly alternative to the use of chemical pesticides. Control of fungal plant diseases using naturally occurring non-pathogenic microorganisms represents a promising approach for the control of plant disease. Biological control using antagonistic microbes alone, or as supplements to minimize the use of chemical pesticides in an integrated plant disease management system, has become more prevalent in recent years.

*Rhizoctonia solani* Kühn [teleomorph: *Thanatephorus cucumeris* (Frank) Donk] is one of the most important, universal, destructive soilborne phytopathogenic fungi that causes various diseases in many plant species world-wide. It is responsible for the damping-off on seedlings and root rot, stem rot, stem canker, foliage blights or spots near the ground and rotting of storage organs (Banville et al., 1996; Boosalis and Scharen, 1959).

Mycoparasites and their possible use in biological control have been recognized since 1932. Since that time there are a number of examples of fungi that parasitize plant pathogens including *R. solani*. Considerable work has focused on various *Trichoderma* and *Gliocladium* species; they have been studied to the greatest extent and possess great potential as biocontrol agents (Weindling, 1932; Barnett and Binder, 1973 ; Lumsden, 1981; Elad et al., 1980, 1983; Papavizas, 1985; Adams, 1990; Coley-Smith et al., 1991; Wilson et al., 1992; Cuevas et al., 1995; Lumsden et al., 1995; Hjeljord and Tronsmo, 1998; Kredics et al., 2003; Samuels, 2006; Aydın and Turhan, 2013).

*Trichoderma* spp. are basically soil-borne saprophytic fungi which possess a symbiotic relationship with the plant rhizosphere. They are well known for their antagonism against several fungi, to produce different antibiotic substances and to inhibit the growth of pathogenic fungi by modifying the rhizosphere. The antagonistic properties of *Trichoderma* spp. towards fungal pathogens are based on strong mycoparasitism, antibiosis, competition for nutrients and space, promotion of plant growth and plant defense responses. They are widely used as commercial biofungicides all over the world (Dennis and Webster, 1971; Chet and Baker, 1981; Chet et al., 1981; Howell, 2003, 2006; Benítez et al., 2004; Harman et al., 2004; Harman, 2006; Almeida et al., 2007; Verma et al., 2007; Woo and Lorito, 2007; Bailey et al., 2008; Vinale et al., 2008; Amin et al., 2010; Anees et al., 2010; Brotman et al., 2010; Chaverri et al., 2015; Hermosa et al., 2012; Kumar et al., 2012; Hyakumachi, 2013; Abo-Elyousr et al., 2014; Asad et al., 2014; Mukherjee et al., 2014; Naher et al., 2014; Rinu et al., 2014).

The isolation of *Trichoderma* and *Gliocladium* from soil was investigated frequently by researchers using a variety of methods such as “serial dilution plate technique”, “pour plate”, “differential centrifugation technique”, “the slide-trap method” and “the dilution plate count technique” (Hopkins et al., 1991; Aydın and Turhan, 2009; Vargas Gil et al. 2009).

Furthermore, several authors have reported the use of agar cultures of fungi or their resting bodies like sclerotia as selective bait for the isolation of specific mycoparasites from soil or from natural habitats. It is possible to specifically select the mycoparasitic fungi by using agar plate precolonized with a suitable host fungus as a bait. Promising results have been obtained using this method and various mycoparasitic fungi including *Trichoderma* spp., *Gliocladium* spp. and others could be isolated (Deacon and Henry, 1978; Foley and Deacon, 1985; van den Boogert and Gams, 1988; Lodha and Webster, 1990; van den Boogert et al., 1990; Deacon and Berry, 1992; Mulligan and Deacon, 1992; Mulligan et al., 1995; Knudsen et al., 1997; Crauss et al., 1998; Molan, 2009).

Ribeiro and Butler (1992) used the “sclerotia bait technique” for the isolation of *Pythium* species mycoparasitic on *Sclerotinia sclerotiorum*. *Trichoderma* strains were isolated from *S. sclerotiorum* sclerotia by selective isolation employing the sclerotial bait technique (Melo et al., 1996). Ali-Shtayeh and Saleh (1999)

reported that the sclerotia baiting technique was highly effective for the detection and isolation of some mycoparasitic *Pythium* species from soils assayed. Baiting and filtering techniques are used to recover species of *Phytophthora* and *Pythium* from water and soil (Roberts et al., 2005). Karaca et al. (2008) used the sclerotia of *Botrytis cinerea* and *S. sclerotiorum* as baits for the isolation of the mycoparasitic fungi from Oomycetes. Burgess and Hepworth (1996) used sclerotia of *S. minor* as baits to isolate *T. virens* from soil. Selective isolation of mycoparasites of *S. minor* from soil by baiting proved to be useful for enumerating *T. virens* on roots of sunflower and showed promise for studies of the ecology of biocontrol agents added to soil or seed (Isnaini et al., 1998). Human hair baiting technique was followed to isolate dermatophytes and related keratinophilic fungi from the soil samples (Kachuei et al., 2012; Malek et al, 2013; Deshmukh and Verekar, 2014). The larvae of *Galleria mellonella*, were used as a standard bait insect for the isolation of entomopathogenic fungi, because the larvae are highly susceptible to infection by these fungi (Chandler et al. 1997; Meyling, 2007; Goble et al., 2010, 2012; Hernández-Domínguez et al., 2016).

The objective of this study was to improve the isolation efficiency of mycoparasitic fungi, especially *Trichoderma* spp. and *Gliocladium* spp. from soil applying precolonized *R. solani* culture as a trap in the first stage and using a modified nutrient medium supplemented with tolclaphos-methyl in the second.

## MATERIALS AND METHODS

### The isolation process of *R. solani* mycoparasites

For the isolation of *Rhizoctonia* mycoparasites, the soil samples were taken from 11 different ecological habitat in Turkey. They were collected from a depth of 1-15 cm, litter-free layer; transported to the laboratory in plastic bags and processed immediately. An isolate of *R. solani* recovered from diseased potato roots was used for all experiments. First, *R. solani* was allowed to cover the PDA plate completely. Then somewhat soil was placed onto the precolonized plate and incubated in the dark at 24 °C for two weeks. After removal of the soil particles, the underlying *Rhizoctonia* colony was thoroughly washed with sterilized water. Potato Dextrose Agar (Merck 1.10130) was sterilized in autoclave after adding 33 mg/L Rose Bengal (BDH Chemicals). Than Streptomycin sulfate (Sigma, 200 mg/L) dissolved in 10 ml sterile distilled water and tolclaphos-methyl (Rizolex®, Sumimoto, 12 mg/L) dissolved in 10 ml 95% ethyl alcohol were added after the sterile medium had cooled to 50°C. Discs (5 mm) cut from the freshly washed *R. solani* colony were transplanted (mycelial side beneath) onto the modified PDA containing rose bengal, streptomycine and tolclaphos-methyl (33 mg/L, 200 mg/L and 6 mg active ingredient/L, respectively) and reincubated approximately for two weeks on a laboratory bench at room temperature (Figure 1, 2). Variegated fungi grown in the meanwhile were isolated into pure cultures and evaluated then for their potential to parasitize *R. solani* in dual cultures.

### Demonstrating the mycoparasitic nature of the isolates

A mycelial disc (5 mm) of the potential mycoparasitic isolate was placed onto the surface of modified PDA (ten times diluted PD broth + 15 g/l agar). According to growth rate of candidate antagonist, either simultaneously or two days later, a mycelial disc of *R. solani* was placed 4 cm apart onto the same plate. The mycelial disc of pathogen or antagonist grown alone served as control. Then the plates were incubated in the dark at 24 °C. As from the paired colonies reached each others after some days, overgrowth of *R. solani* by mycoparasitic isolate could be detected. Directly interaction area or only mycelial samples taken from there were examined under light microscope. Throughout the experiments, there were three replications.

## RESULTS AND DISCUSSION

When the colonies grown in dual cultures come into contact, the mycoparasite has continued to grow rapidly through the *Rhizoctonia* colonies. When cultures intermingled, the physical interaction between the mycoparasites and *R. solani* was examined by light microscopy. Hyphae of *R. solani* and hyperparasites were distinguished from each other by the orientation of hyphal growth. The fine hyphae of the mycoparasites were also readily

A NOVEL TECHNIQUE FOR THE RECOVERY, ISOLATION AND PRELIMINARY EVALUATION OF  
*RHIZOCTONIA SOLANI* MYCOPARASITES FROM SOIL

differentiated from the coarse hyphae of *Rhizoctonia*. Tight coiling by the mycoparasites around *Rhizoctonia* hyphae was observed. The internal parasitism was evidenced by the fact that parasitic hyphae were frequently found inside the host hyphae



**Figure 1.** Mature *Rhizoctonia solani* colony on PDA (a), waiting period as covered with soil sample (b)



**Figure 2.** Discs cut from thoroughly washed colony (c), rapid growth of *Rhizoctonia solani* alone on PDA having Rose Bengal and Streptomycin (d), growth of mycoparasites on the same medium having additionally tolclaphos-methyl (e).

By using the novel technique introduced in this study, 264 mycoparasitic isolates were obtained from soil. The majority of the isolates were found to be in the genera of *Trichoderma* and *Gliocladium*. Fourteen *Trichoderma* species, *Gliocladium roseum* Bainier, *Chaetomium* sp., *Geotrichum* sp., *Paecilomyces* sp., *Papulospora* sp. and a *Spicaria* sp. were determined as having mycoparasitic activity against *R. solani* as well. Identification of mycoparasites made by Prof. Dr. W. Gams, Prof. Dr. G. J. Samuels and Prof. Dr. G. Turhan.

Mycoparasitism, the direct attack of one fungus to another, is a very complex process. It has been known for many years that the genus *Trichoderma* has a strong mycoparasitic activity; it can sense the presence of target fungi;

it is attracted to grows towards its host and appeared to grow typically, probably by chemotropism toward them (Chet et al., 1981; Hyakumachi, 2013).

*Trichoderma harzianum*, *T. viride*, *T. pseudokoningii*, *T. hamatum*, *T. virens*, *T. koningii*, *T. aureoviride* and *Trichoderma* spp. were previously isolated by using known conventional methods in Turkey (Turhan, 1973; İren et al., 1988; Çeliker and Nemli, 1994).

It is possible to specifically select the mycoparasitic fungi by using precolonized agar plate with the host fungus. Only fungi that are able to derive nutrients from the precolonizing fungus and to tolerate its waste products are able to grow. Unfortunately there is no universally susceptible host fungus that can be used in this technique. Since all mycoparasitic fungi have restricted host ranges, the host fungus used to precolonize the plate will greatly influence the type of mycoparasite that will be detected. Mycoparasites isolated by this method include *Gliocladium roseum* and related species (Foley and Deacon, 1985), *Papulaspora* sp. (Deacon and Berry, 1992; Mulligan and Deacon, 1992), *Pythium oligandrum* and related species (Deacon and Henry, 1978; Foley and Deacon, 1985; Mulligan et al., 1995) and *Verticillium biguttatum* (van den Boogert and Gams, 1988).

In the present study the precolonized plate method was used at the first stage of the technique. But, because of the extremely fast growing rate of *R. solani*, it was difficult to catch it by invaders, even though it is already parasitized during the waiting period beneath the soil coating in petri dishes. Hence *R. solani* grew first from the precolonized discs and completely filled the plate in a little while (Figure 2d) unless the presence an intentional inhibitive in the culture medium.

The results of a study by Hameed (2008) showed that Rizolex® (50% Tolclofos methyl) was effective on radial growth of *R. solani*. There were differences between *R. solani* and *T. harzianum* in response to Rizolex, then isolates of *R. solani* were inhibited completely at all concentrations tested, while *T. harzianum* was affected lesser. Yobo et al.(2010) found that although *R. solani* growth was completely inhibited in concentrations of 0.05 and 0.125 g ai/l tolclofos-methyl, *Trichoderma* isolates tested were less sensitive to this chemical.

We hypothesized that *R. solani* is more susceptible to tolclofos-methyl than *Trichoderma* and *Gliocladium* spp. Based on this hypothesis, we used a special growth medium supplemented with tolclofos methyl. The purpose of adding this chemical into the isolation medium was to prevent extreme speedy growth of *R. solani*, which already existed on precolonized discs. Really, the presence of tolclofos-methyl in nutrient medium prevented the unwelcome growth of *R. solani* significantly. Fortunately, *Trichoderma* and *Gliocladium* isolates were least affected displaying considerable tolerance to this fungicide and they could grow well on this modified PDA medium.

On account of the lesser sensitivity of two foremost important mycoparasitic genera to tolclofos-methyl than *R. solani*, it can be concluded that this chemical could be used in integrated management in addition to biocontrol agents against this pathogen.

Furthermore, the technique described, without doubt, may also be used for detecting and isolating the mycoparasites of other fast growing fungus and fungus-like organisms such as *Pythium* from soil by changing the supplemented chemical which suppress only the target pathogen.

#### ACKNOWLEDGEMENT

Identification of mycoparasites by Prof. Dr. W. Gams and Prof. Dr. G. J. Samuels is gratefully acknowledged.

#### LITERATURE CITED

- Abo-Elyousr, K.A.M., Abdel-Hafez, S.I.I. and Abdel-Rahim. I.R. 2014. Isolation of *Trichoderma* and evaluation of their antagonistic potential against *Alternaria porri*. Journal Phytopathology 162: 567–574
- Adams, P.B. 1990. The Potential of Mycoparasites for Biological Control of Plant Diseases. Annual Review Phytopathology 28: 59-72.
- Ali-Shtayeh, M.S. and Saleh, A.S.F. 1999. Isolation of *Pythium acanthicum*, *P. oligandrum*, and *P. periplocum* from soil and evaluation of their mycoparasitic activity and biocontrol efficacy against selected phytopathogenic *Pythium* species. Mycopathologia 145: 143–153.

A NOVEL TECHNIQUE FOR THE RECOVERY, ISOLATION AND PRELIMINARY EVALUATION OF  
*RHIZOCTONIA SOLANI* MYCOPARASITES FROM SOIL

- Almeida, F., Cerqueira, F., Silva, R., Ulhoa, C. and Lima A. 2007. Mycoparasitism studies of *Trichoderma harzianum* strains against *Rhizoctonia solani*: evaluation of coiling and hydrolytic enzyme production. *Biotechnology Letter* 29: 1189–1193.
- Amin, F., Razdan, V.K., Mohiddin, F.A., Bhat, K.A. and Banday, S. 2010. Potential of *Trichoderma* species as biocontrol agents of soil borne fungal propagules. *Journal of Phytology* 2: 38–41.
- Anees, M., Tronsmo, A., Edel-Hermann, V., Hjeljord, L.G., Heraud, C., and Steinberg, C. 2010. Characterization of field isolates of *Trichoderma* antagonistic against *Rhizoctonia solani*. *Fungal Biology* 114: 691-701.
- Asad, S.A., Ali, N., Hameed, A., Khan, S.A., Ahmad, R., Bilal, M., Shahzad, M. and Tabassum, A. 2014. Biocontrol efficacy of different isolates of *Trichoderma* against soil borne pathogen *Rhizoctonia solani*. *Polish Journal of Microbiology* 63: 95–103.
- Aydın, M.H., Turhan, G. 2009. *Rhizoctonia solani*'nin Fungal Antagonistlerinin Belirlenmesi Üzerinde Araştırmalar. "Studies on determination of fungal antagonists of *Rhizoctonia solani*". *Anadolu Journal of AARI* 19: 49-72.
- Aydın, M.H., Turhan, G. 2013. Patateste *Rhizoctonia solani*' ye Karşı *Trichoderma* Türlerinin Etkinliği ve Bazı Fungisitlerle Birlikte Kullanılması. "The efficacy of *Trichoderma* species against *Rhizoctonia solani* on potato and their Integration with some fungicides". *Anadolu Journal of AARI* 23: 12-30.
- Bailey, B., Bae, H., Strem, M.D., Crozier, J., Thomas, S.E., Samuels, G.J., Vinyard, B.T. and Holmes, K.A. 2008. Antibiosis, mycoparasitism, and colonization success for endophytic *Trichoderma* isolates with biological control potential in *Theobroma cacao*. *Biological Control* 46: 24–35.
- Banville, J.B., Carling, E.C. and Otrysko, B.E.1996. *Rhizoctonia* Diseases on Potato. pp. 321-330, in *Rhizoctonia* Species: Taxonomy, Molecular biology, Ecology, Pathology and Disease Control, Edited by. B. Sneh., S. Jabaji-Hare., S. Neate and G. Dijst. Kluwer Academic Publishers, London.
- Barnett, I.I.L., Binder, F.L.1973. The fungal host-parasite relationship. *Annual Review Phytopathology* 11: 273-292
- Benítez, T., Rincón, A.M., Limón, M.C., Codón, A.C. 2004. Biocontrol mechanisms of *Trichoderma* strains. *International Microbiology* 7: 249–260.
- Brotman, Y., Kapuganti, J.G., Viterbo, A. 2010. *Trichoderma*. *Current Biology* 20, R390-R391.
- Burgess, D.R., Hepworth, G. 1996. Biocontrol of sclerotinia stem rot (*Sclerotinia minor*) in sunflower by seed treatment with *Gliocladium virens*. *Plant Pathology* 45: 583-592.
- Boosalis, M.G. and Scharen, A.L 1959. Methods for microscopic detection of *Aphaanomyces euteiches* and *Rhizoctonia solani* and for isolation of *Rhizoctonia solani* associated with plant debris. *Phytopathology*, 49: 192-198.
- Chandler, D., Hay, D., Reid, A.P. 1997. Sampling and occurrence of entomopathogenic fungi and nematodes in UK soils. *Applied Soil Ecology*. 5: 133–141.
- Chaverri, P., Branco-Rocha, F., Jaklitsch, W., Gazis, R., Degenkolb, T., Samuels, G.J.2015. Systematics of the *Trichoderma harzianum* species complex and the re-identification of commercial biocontrol strains. *Mycologia*, 107: 558–590.
- Chet, I., Harman, G.E., Baker, R. 1981. *Trichoderma hamatum*: Its hyphal interactions with *Rhizoctonia solani* and *Pythium* spp. *Microbiological Ecology* 7: 29-38.
- Chet, I. 1987. *Trichoderma*: Application, Mode of Action and Potential as a Bio Control Agent of Soil Borne Plant Pathogenic Fungi. In: Chet, I. (Ed.), *Innovative Approaches to Plant Disease Control*. Wiley, New York, pp. 137-160.
- Chet, I., Baker, R. 1981. Isolation and biocontrol potential of *Trichoderma hamatum* from soil naturally suppressive to *Rhizoctonia solani*. *Phytopathology* 71: 286-290.

- Coley-Smith, J.R., Ridout, C.T., Mitchell, C.M., Lynch, J.M. 1991. Control of bottom rot disease of lettuce (*Rhizoctonia solani*) using preparations of *Trichoderma viride*, *T. harzianum* or tolclofos-methyl. *Plant Pathology*. 40: 359-366.
- Crauss, U., Bidwell, R., Ince, J. 1998. Isolation and Preliminary Evaluation of Mycoparasites as Biocontrol Agents of Crown Rot of Banana. *Biological. Control* 13: 111-119.
- Cuevas, V.C., Soriano, J.M., Bagunu, L.G., Soniega, J.A., Alfonso, A.L. 1995. Control of damping-off diseases of vegetables by *Trichoderma* species. *Philippine Agriculturist* 78, 255-276.
- Çeliker, N.M. and Nemli, T. 1994. Investigation on biocontrol of white root rot [*Rosellinia necatrix* (Hartig) Berlese]. *Türkiye III. Biyolojik Mücadele Kongresi* 25-28 Ocak.
- Deacon, J.W. and Berr, A. 1992. Modes of action of mycoparasites in relation to biocontrol of soilborne plant pathogens. In: Tjamos, E.C. and Cook, R.J., (Ed.), *Biological control of plant diseases*. Plenum Press, New York, N.Y, pp. 157-165.
- Deacon, J.W., Henry, C.M. 1978. Mycoparasitism by *Pythium oligandrum* and *P. acanthicum*. *Soil Biology and Biochemistry* 10: 409-415.
- Dennis, C., Webster, J. 1971. Antagonistic properties of species-groups of *Trichoderma*: III. Hyphal interaction. *Transactions the British. Mycological. Society* 57: 363-369.
- Deshmukh, S.K., Verekar, S.A. 2014. Isolation of keratinophilic fungi from selected soils of Sanjay Gandhi National Park, Mumbai (India). *Journal Medical Mycology* 24: 319-327.
- Elad, Y., Barak, R., Chet, I. and Heris, Y. 1983. Ultrastructural studies of the interaction between *Trichoderma spp.* and plant pathogenic fungi, *Phytopathology* 107: 168-175.
- Elad, Y., Chet, I. and Katan, J. 1980. *Trichoderma harzianum* a biocontrol agent effective against *Sclerotium rolfsii* and *Rhizoctonia solani*. *Phytopathology* 70: 119-121.
- Foley, M.F. and Deacon, J.W. 1985. Isolation of *Pythium oligandrum* and other necrotrophic mycoparasities from soil. *Transactions British Mycological Society* 85: 631-639.
- Goble, T.A., Dames, J.F., Hill, M.P., Moore, S.D. 2010. The effects of farming system, habitat type and bait type on the isolation of entomopathogenic fungi from citrus soils in the Eastern Cape Province, South Africa. *Biocontrol* 55: 399-412.
- Goble, T.A., Costet, L., Robene, I., Nibouche, S., Rutherford, R.S., Conlong, D.E., Hill, M.P. 2012. *Beauveria brongniartii* on white grubs attacking sugarcane in South Africa. *Journal of Invertebrate Pathology*, 111: 225-236.
- Hameed, F.R. 2008. Effect of Rizolex on *Rhizoctonia solani* Kühn isolates and biocontrol fungus *Trichoderma harzianum* Rifai *in vitro*. *Journal of Kerbala University* 4: 218-223.
- Harman, G.E. 2006. Overview of mechanisms and uses of *Trichoderma spp.* *Phytopathology* 96: 190-194.
- Harman, G.E., Howell, C.R., Viterbo, A., Chet, I., Lorito, M. 2004. *Trichoderma* species: opportunistic, avirulent plant symbionts. *Nature Reviews Microbiology* 2: 43-56.
- Hermosa, R., Viterbo, A., Chet, I., Monte, E. 2012. Plant beneficial effects of *Trichoderma* and its genes. *Microbiology*, 158: 17-25.
- Hernández-Domínguez, C., María de Lourdes C.B, M.L., Alvarado-Aragón, L.U., Hernández-López, G., Guzmán-Franc, A.W. 2016. Comparison of the relative efficacy of an insect baiting method and selective media for diversity studies of *Metarhizium* species in the soil. *Biocontrol Science Technology* 26: 707-717.
- Hjeljord, L., Tronsmo, A. 1998. *Trichoderma* and *Gliocladium* in biological control: an overview: In: Harman, G.E., Kubicek, C.P. (Eds.) *Trichoderma and Gliocladium: Enzymes, Biological control and Commercial Applications*. Taylor and Francis Ltd., London, pp. 131-151.
- Hopkins, D.W., Macnaughton, S.J., O' Donnell, A.G. 1991. A dispersion and differential centrifugation technique for representatively sampling microorganisms from soil. *Soil Biology Biochemistry* 23: 217-225.

A NOVEL TECHNIQUE FOR THE RECOVERY, ISOLATION AND PRELIMINARY EVALUATION OF  
*RHIZOCTONIA SOLANI* MYCOPARASITES FROM SOIL

- Howell, C.R. 2003. Mechanisms employed by *Trichoderma* species in the biological control of plant diseases: The history and evolution of current concepts. *Plant Diseases* 87: 4–10.
- Howell, C.R. 2006. Understanding the mechanisms employed by *Trichoderma virens* to effect biological control of cotton diseases. *Phytopathology* 96: 178–180.
- Hyakumachi, M. 2013. Research on biological control of plant diseases: present state and perspectives. *J. Gen. Plant Pathology* 79: 435–440.
- Isnaini, M., Burgess, D.R., Keane, P.J. 1998. The use of cultures of *Sclerotinia minor* for selective isolation and enumeration of mycoparasitic isolates of *Trichoderma* from soil and roots. *Australasian Plant Pathology*. 27: 244-250.
- İren, S., Maden, S., Katırcıoğlu, Z., Erzurum, K. 1988. *Trichoderma* species determined in Turkey. *Journal of Turkish Phytopathology* 17: 107.
- Kachuei, R., Emami, M., Naeimi, B., Diba, K. 2012. Isolation of keratinophilic fungi from soil in Isfahan province, *Iranian Journal of Medical Microbiology* 22: 8-13.
- Karaca, G., Tepedelen, G., Belghouthi, A., Paul, B. 2008. A new mycoparasite, *Pythium lycopersicum*, isolated in Isparta, Turkey: Morphology, molecular characteristics and its antagonism with phytopathogenic fungi. *FEMS Microbiology. Letters* 288, 163–170.
- Knudsen, I.M.B., Hockenhull, J., Funck, J.D., Gerhardson, B., Hökeber, M., Tahvonen, R., Teperi, E., Sundheim, L., Henriksen, B. 1997. Selection of biological control agents for controlling soil and seed-borne diseases in the field. *European Journal of. Plant Pathology* 103: 775–784.
- Kredics, L., Antal, Z., Manczinger, L., Szekeres, A., Kevei, F., Nagy, E. 2003. *Trichoderma* strains with biocontrol potential, *Food Technology Biotechnology* 41: 37-42.
- Kumar, K., Amaresan, N., Bhagat, S., Madhuri, K., Srivastava, R.C. 2012. Isolation and Characterization of *Trichoderma* spp. for Antagonistic Activity Against Root Rot and Foliar Pathogens. *Indian Journal of. Microbiology* 52: 137–144.
- Lodha, B.C., Webster, J. 1990. *Pythium acanthophoron*, a mycoparasite, rediscovered in India and Britain. *Mycology Research*. 94: 1006–1008.
- Lumsden, R.D. 1981. Hyperparasitism for control of plant pathogens. In: Pimentel, D. (Ed.), *CRC Handbook of Pest Management in Agriculture*, Vol. 1, CRC Press, Boca Raton, Florida. pp. 475-484.
- Lumsden, R.D., Lewis, J.A., Ravel, D.R. 1995. Formulation and delivery of biocontrol agents for use against soilborne plant pathogens. In: Hall, F.R., Bary, J.W. (Eds.), *Biorational Pest Control Agents, Formulation and Delivery*, ACS Symposium Series 595, Washington, pp. 166-182.
- Malek, E., Moosazadeh, M., Hanafi, P., Nejat, Z.A., Amini, A., Mohammadi, R., Kohsar, F., Niknejad, F. 2013. Isolation of Keratinophilic Fungi and Aerobic *Actinomycetes* From Park Soils in Gorgan, North of Iran, *Jundishapur Journal of Microbiology* 6 :10, e11250.
- Melo, I.S., Faull, J.L., Graeme-Cook, K.A. 1996. *Trichoderma koningii* and *Trichoderma harzianum* as destructive mycoparasites of *Sclerotinia sclerotiorum*. In: *International Congress of Mycology Division, 8, Proceedings*. Jerusalém, 1996, p.183.
- Meyling, N. 2007. Methods for isolating entomopathogenic fungi from the soil environment. *Laboratory Manual 2007*, Department of Ecology, Faculty of Life Sciences, University of Copenhagen, Denmark.
- Molan, Y.Y. 2009. Detection of Presumptive Mycoparasites in Soil Placed on Host-Colonized Agar Plates in Riyadh Region, Saudi Arabia. *Asian Journal of Plant Pathology* 3, 22-26.
- Mukherjee, A.K., Kumar, A.S., Kranthi, S., Mukherjee, P.K. 2014. Biocontrol potential of three novel *Trichoderma* strains: isolation, evaluation and formulation. *3 Biotechnique*. 4, 274-281
- Mullingan, D.F.C., Deacon, J.W. 1992. Detection of presumptive mycoparasites in soil placed on host-colonized agar plates. *Mycology Research* 96, 605-608.



- Mulligan, D.F.C., Jones, E.E., Deacon, J.W. 1995. Monitoring and manipulation of populations of *Pythium oligandrum*, *Pythium mycoparaticum* and a *Papulaspora* species in soil. *Soil Biology Biochemistry* 27, 1333–1343.
- Naher, L., Yusuf, U.K., Ismail, A. and Hossain, K. 2014. *Trichoderma* spp.: A Biocontrol Agent for Sustainable Management of Plant Diseases. *Pakistan Journal of Botany* 46, 1489-1493.
- Pal, K. K. and Gardener, B.M.S. 2006. Biological Control of Plant Pathogens. The Plant Health Instructor DOI: 10.1094/PHI-A-2006-1117-02.
- Papavizas, G.C. 1985. *Trichoderma* and *Gliocladium*: biology, ecology, and potential for biocontrol. *Annual Review Phytopathology* 23, 23–54.
- Ribeiro, W.R.C. and Butler, E.E. 1992. Isolation of mycoparasitic species of *Pythium* with spiny oogonia from soil in California. *Mycology Research* 96, 857–862.
- Rinu, K., Sati, P., Pandey, A. 2014. *Trichoderma gamsii* (NFCCI 2177): A newly isolated endophytic, psychrotolerant, plant growth promoting, and antagonistic fungal strain. *Journal of Basic Microbiology* 54, 408–417.
- Roberts, P.D., Urs, R.R., French-Monar, R.D., Hoffine, M.S., Seijo, T.E., Mcgovern, R.J. 2005. Survival and recovery of *Phytophthora capsici* and oomycetes in tailwater and soil from vegetable fields in Florida. *Annals of Applied Biology* 146, 351–359.
- Samuels, G.J. 2006. *Trichoderma*: Systematics, the sexual state, and ecology. *Phytopathology* 96, 195-206.
- Turhan, G. 1973. Fungi isolated from the roots of diseased vegetable seedlings. *Journal of Turkish Phytopathology* 2, 100-112.
- van den Boogert, P.H.J.F. and Gams, W. 1988. Assessment of *Verticillium biguttatum* in agricultural soils. *Soil Biology and Biochemistry*. 20, 899-905.
- van den Boogert, P.H.J.F., Jager, G. and Velvis, H. 1990. *Verticillium biguttatum*, an important mycoparasite for the control of *Rhizoctonia solani* in potato. In: Hornby, D. (Ed.), *Biological Control of Soil-borne Plant Pathogens*. CAB International, Wallingford. pp. 77-91.
- Vargas Gil, S., Pastor, S., March, G.J. 2009. Quantitative isolation of biocontrol agents *Trichoderma* spp., *Gliocladium* spp. and actinomycetes from soil with culture media. *Microbiology Research* 164, 196–205.
- Verma, M., Brar, S., Tyagi, R., Surampalli, R. and Valero, J. 2007. Antagonistic fungi, *Trichoderma* spp.: panoply of biological control. *Biochemical Engineering Journal* 37, 1–20
- Vinale, F., Sivasithamparam, K., Ghisalberti, S., Lorito, M. 2008. *Trichoderma*–plant– pathogen interactions. *Soil Biology and Biochemistry* 40, 1–10.
- Weindling, R. 1932. *Trichoderma lignorum* as a parasite of other soil fungi. *Phytopathology* 22, 837-45.
- Wilson, M., Crawford, E.K., Campbell, R. 1992. Biological control by *Trichoderma harzianum* of damping-off of lettuce caused by *Rhizoctonia solani*. *Bulletin EPPO* 18, 83-89.
- Woo, S.L., Lorito, M. 2007. Exploiting the Interactions Between Fungal Antagonists, Pathogens and the Plan For Biocontrol. In: Vurro, M., Gressel, J. (Eds.), *Novel Biotechnologies for Biocontrol Agent Enhancement and Management*, Springer, Dordrecht, NL., pp. 107–130.
- Yobo, K. S., Laing, M. D., Hunter, C. H. 2010. Application of selected biological control agents in conjunction with tolclofos-methyl for the control of damping-off caused by *Rhizoctonia solani*. *African Journal of Biotechnology* 9, 1789-1796.

